

1. About the data set

Site name (AsiaFlux three letter code)	Kahoku Experimental Watershed (KHW)	
Period of registered data	From January 1, 2007 to December 31, 2007	
This document file name	KHW_2007_001a.pdf	
Corresponding data file name	KHW_2007_001.csv	
Revision information		
Date	Details of revision	Renewed file name
31 July 2014	First registration	KHW_2007_001a.pdf KHW_2007_001.csv
Contact person#1	[Flux&Meteorology] Takanori SHIMIZU (simizuta@ffpri.affrc.go.jp)	
Contact person#2		
Contact person#3		
Contact person#4		

2. Site description

Hour line (Time difference from UTC)	Japan Standard Time (JST) (9 hours ahead of UTC)
Location (address)	Iwano, Kahoku-town, Yamaga-city, Kumamoto pref., Japan
Position	33.137N, 130.7095E (World Geodetic System 1984, GPS: Garmin eTrex Legend and map)
Elevation	165m above sea level (World Geodetic System 1984, GPS: Garmin eTrex Legend and map)
Terrain type	Rolling terrain
Slope	16 degrees (average value around the tower)
Area	approx. 12.7ha
Fetch	>300m
Climate	Warm temperate (Köppen climate classification: Cfa)
Mean annual air temperature	15.3 degrees C (2000-2008)
Mean annual precipitation	2138mm (2000-2008)
Vegetation Type	Evergreen coniferous forest
Dominant Species (Overstory)	<i>Cryptomeria japonica</i> (Sugi cedar), <i>Chamaecyparis obtuse</i> (Hinoki cypress): plantimal <i>Castanopsis cuspidata</i> , <i>Fagus japonica</i> (japanese beech): grown in gaps
Dominant Species (Understory)	<i>Fagus japonica</i> Maxim., <i>Castanopsis sieboldii</i> , etc.
Canopy height	10-35m
Breast high diameter	50cm (max, Sugi cedar)
Age	Around 50 years (Sugi cedar), 30-60 years (Hinoki cypress)
LAI	3.6-5.2 (estimated by LAI-2000)
Soil Type	Brown forest soil B _D
Other information	

Reference

--

4. Registered Data

Observation items	Symbol	Unit	Height(s) Depth(s)	Instruments	Note
Date	DATE	-	-	-	yyyymmdd
Time	TIME	-	-	-	hhmm
Precipitation	PPT	mm	1.5m	RT-5 (IKEDA)	See Note [1]
Air temperature	Ta	degrees C	41.5m	ML-020L (EKO)	See Note [2]
Relative humidity	Rh	%	41.5m	ML-202L (EKO)	See Note[2], [3]
Wind speed	U	$\text{m}\cdot\text{s}^{-1}$	51.0m	DAT-600-3T (KAIJO)	
Wind direction	WD	degrees	51.0m	DAT-600-3T (KAIJO)	
Global solar radiation (incoming / downward)	Sd	$\text{W}\cdot\text{m}^{-2}$	47.2m	CM14 (Kipp & Zonen)	See Note [4]
Reflected solar radiation (upward)	Su	$\text{W}\cdot\text{m}^{-2}$	47.2m	CM14 (Kipp & Zonen)	See Note [4]
Photosynthetic active photon flux density (downward)	Pd	NA	NA	NA	
Reflected PAR (upward)	Pu	NA	NA	NA	
Net radiation	Rn	$\text{W}\cdot\text{m}^{-2}$	47.2m	CM14 (Kipp & Zonen) CG3 (Kipp & Zonen)	See Note [5], [6]
Soil heat flux	G	$\text{W}\cdot\text{m}^{-2}$	-0.05m	HFT3.1 (REBS)	
Sensible heat flux	H	$\text{W}\cdot\text{m}^{-2}$	51.0m	DAT-600-3T (KAIJO)	
Latent heat flux	IE	$\text{W}\cdot\text{m}^{-2}$	51.0m	DAT-600-3T(KAIJO) LI-7000 (LI-COR)	Closed-path system QC
Friction velocity	Ust	NA			
CO ₂ flux	Fc	NA			
Storage change in canopy air layer	Sc	NA			
Net ecosystem exchange	NEE	NA			
Ecosystem respiration	Re	NA			
Gross primary production	GPP	NA			

Note

- [1] Data gaps were filled with data obtained using AMeDAS (weather station managed by Meteorological Agency) data.
- [2] Data gaps were filled with data obtained using the backup system. When data of both observation sets were unavailable, linear interpolation were applied. The residual gaps were filled with AMeDAS data.
- [3] value of >100% is replaced by 100%
- [4] value in night time is replaced by 0.0.

[5] $R_n = S_d - S_u + L_d - L_u$ (Ld: downward longwave radiation, Lu: upward longwave radiation_)

[6] Data gaps were filled with data obtained by the net radiometer; NR-LITE (Kipp&Zonen).

Reference

Data format

Data consists of fixed length (8 digits) comma separated format. Missing data is labeled as "-9999.00"

Line 1: Symbol (Date, Time, PPT, Ta,

Line 2: Unit (yyyymmdd, hhmm, mm, degC,

"hhmm" shows intermediate time of averaging period.

i.e. "1215" labels half-hourly average (or sum) of data from 12:00 to 12:30

Line 3: Comment

Line 4: Data

:

Data Example

Date, Time, yyyymmdd, hhmm,	PPT, mm,	Ta, degC,	Rh, %,	U, ms ⁻¹ ,	WD, deg,	Sd, Wm ⁻² ,	Su, Wm ⁻² ,	Pd, (#1),	Pu, (#1),	
File= KWG_2000_001.CSV; Created: 20100326; Gap= -9999.0; (*1): micro-mol m ⁻² s ⁻¹										
20000101, 0015,	0.0,	3.34,	87.19,	1.58, -9999.0,	0.1, -9999.0,	0.1, -9999.0,	0.1, -9999.0,	0.1,	0.0,	
20000101, 0045,	0.0,	3.12,	88.14,	1.44, -9999.0,	0.0, -9999.0,	0.0, -9999.0,	0.1, -9999.0,	0.1,	0.0,	
20000101, 0115,	0.0,	2.36,	90.51,	1.15, -9999.0,	-0.3, -9999.0,	0.1, -9999.0,	0.1, -9999.0,	0.1,	0.0,	
20000101, 0145,	0.0,	2.14,	91.32,	0.83, -9999.0,	0.0, -9999.0,	0.0, -9999.0,	0.1, -9999.0,	0.1,	0.0,	
20000101, 0215,	0.0,	2.28,	88.96,	0.49, -9999.0,	-0.3, -9999.0,	-0.3, -9999.0,	0.1, -9999.0,	0.1,	0.0,	
20000101, 0245,	0.0,	2.24,	89.82,	0.35, -9999.0,	-0.2, -9999.0,	-0.2, -9999.0,	0.2, -9999.0,	0.2,	0.0,	
20000101, 0315,	0.0,	2.05,	89.49,	1.50, -9999.0,	0.1, -9999.0,	0.1, -9999.0,	0.2, -9999.0,	0.2,	-0.1,	
20000101, 0345,	0.0,	2.41,	87.25,	1.27, -9999.0,	0.0, -9999.0,	0.0, -9999.0,	0.2, -9999.0,	0.2,	0.0,	
20000101, 0415,	0.0,	2.31,	86.83,	1.12, -9999.0,	-0.2, -9999.0,	-0.2, -9999.0,	0.1, -9999.0,	0.1,	0.0,	
20000101, 0445,	0.0,	2.84,	83.36,	0.54, -9999.0,	-0.6, -9999.0,	-0.6, -9999.0,	0.0, -9999.0,	0.0,	0.0,	
20000101, 0515,	0.0,	2.53,	83.32,	1.23, -9999.0,	0.2, -9999.0,	0.2, -9999.0,	0.2, -9999.0,	0.2,	0.0,	
20000101, 0545,	0.0,	1.59,	87.54,	1.29, -9999.0,	-0.6, -9999.0,	-0.6, -9999.0,	0.0, -9999.0,	0.0,	0.0,	
20000101, 0615,	0.0,	1.89,	85.13,	0.94, -9999.0,	0.4, -9999.0,	0.4, -9999.0,	0.3, -9999.0,	0.3,	0.0,	
20000101, 0645,	0.0,	1.77,	82.40,	0.83, -9999.0,	3.5, -9999.0,	3.5, -9999.0,	8.5, -9999.0,	8.5,	0.4,	
20000101, 0715,	0.0,	2.67,	76.83,	1.38, -9999.0,	45.8, -9999.0,	45.8, -9999.0,	71.9, -9999.0,	71.9,	11.6,	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	

4. Observation and calculation

4-1. Flux observation system and data acquisition

Type of sonic anemometer	DA-600-3T (KAIJO)
Type of IRGA	LI-7000 (LI-COR)
Sampling rate	10Hz
Averaging time	0
Flux measurement height #1	51.0m
Zero-plane displacement	27.0m for southern direction from the tower
Roughness length	5.3m for southern direction from the tower
Calibration information	—
Other information	—

4-2. Flux calculation

Calculation methods		Note
Flow attenuation ^{*5-7}	Transducer shadow correction	Shimizu et al. (1999) Boundary-Layer Met., 93, 227-236.
Coordinate rotation ^{*1-4}	Pitch angle rotation by the ratio of vertical/horizontal wind components moving average ^{*4}	
Lag removal ^{*2, 8, 9}	Automatic	

4-3. Flux corrections

Correction methods		Target flux	Note
Cross wind correction ^{*10, 11}		For sensible heat flux	
Vapor correction			
High frequency loss ^{*4, *13-19}	Combination of theoretical approaches ^{*18}	For latent heat flux (IE)	
	Closed-path H ₂ O correction ^{*4, *19}		
Low frequency loss (Detrending)	Block average ^{*17}	For all	
WPL Correction ^{*20-25}		For Latent heat flux (IE)	
Others ^{*26-28}	Temperature dependency for latent heat: l Humidity dependency for specific heat: Cp Temperature dependency for air density Pressure dependency for air density		

4-4. Quality control ^{*28-29}

QC methods			Note
Raw data test ^{*28,29}	Spike test ^{*27}	Applied	
	Absolute limits	Applied	
	Absolute variance	Applied	
	Higher-moment statistics	skewness	Applied
		kurtosis	Applied
	Discontinuities	Harr mean test	Applied
		Harr variance test	Applied
	Visual inspection	Applied	
Non steady state test ^{*28}		Not applied	
Absolute thresholds			
Others			

4-5. Storage term

Target storage		Note
CO ₂	From CO ₂ profile data (6.0, 11.5, 17.3, 24.0, 31.2, 36.5, 40.3, 46.1m) Sampling interval: 120 seconds at each height	

References

Shimizu, T., Suzuki M. and Shimizu, A. (1999) Examination of a Correction Procedure for the Flow Attenuation in Orthogonal Sonic Anemometers. Boudary-Layer Meteorology, 93(2), 227-236.
--

5. Important events

Date	Events

6. Publications relating to this site

SHIMIZU Takanori (2007) Practical applicability of high frequency correction theories to CO₂ flux measured by a closed-path system. *Boundary-Layer Meteorology*, 122(2):417-438

Publication list: http://www2.ffpri.affrc.go.jp/labs/flux/paper_e.html [KHW]

References

Flux calculation

- *1 McMillen, R.T., 1988. Boundary-Layer Meteorology, 43: 231-245.
- *2 Aubinet M. et al., 2000. Advances in Ecological Research, 30: 113-175.
- *3 Wilczak, J.M., Oncley, S.P. and Stage, S.A., 2001. Boundary-Layer Meteorology, 99: 127-150.
- *4 Shimizu T., 2014. (submitting to Boundary-Layer Meteorology)
- *5 Wyngaard, J. C. and Zhang, S. F., 1985. J. Atmos. Oceanic Tech., 2: 548-558.
- *6 Kaimal, J.C. et al., 1990. Boundary-Layer Meteorol., 53: 103-115.
- *7 Shimizu, T. et al., 1999. Boundary-Layer Meteorol., 64: 227-236.
- *8 Leuning, R. and Judd M.J., 1996. Global Change Biology, 2: 241-254.
- *9 Information from Li-Cor

Flux correction

- *10 Schotanus, P. et al., 1983. Boundary-Layer Meteorology, 26: 81-93.
- *11 Liu, H., Peters, G. and Foken, T., 2001. Boundary-Layer Meteorology, 100: 459-468.
- *12 Kaimal J.C. and Gaynor, J.E., 1991. Boundary-Layer Meteorology, 56: 401-410.
- *13 Watanabe et al., 2000. Boundary-Layer Meteorol. 96, 743-491.
- *14 Massman, W. J., 2000. Agric. For. Meteorol. 104, 185-198
- *15 Massman, W. J., 2001. Agric. For. Meteorol. 107, 247-251
- *16 Moore, C.J., 1986. Boundary-Layer Meteorology, 37: 17-35.
- *17 Moncrieff, J. et al., 2004. Averaging, detrending and filtering of eddy covariance time series. In: X. Lee (Editor), Handbook of Micrometeorology: A guide for surface Flux Measurements. Kluwer, Dordrecht, pp. 7-31.
- *18 Shimizu, T. 2007. Boundary-Layer Meteorology, 122(2):417-438.
- *19 Ibrom, A., et al. 2007. Agricultural and Forest Meteorology, 147:140-156.
- *20 Webb, E. K., Pearman, G.I. and Leuning, R., 1980. Quarterly Journal of the Royal Meteorological Society, 106: 85-100.
- *21 Fuehrer, P.L. and Friehe, C.A., 2002. Boundary-Layer Meteorology, 102: 415-457.
- *22 Liebethal, C. and Foken, T., 2003. Boundary-Layer Meteorology, 109: 99-106.
- *23 Leuning, R. 2004. Measurements of trace gas fluxes in the atmosphere using eddy covariance: WPL corrections revisited. In: X. Lee (Editor), Handbook of Micrometeorology: A guide for surface Flux Measurements. Kluwer, Dordrecht, pp. 119-132.
- *24 Massman, W. 2004. Concerning the measurement of atmospheric trace gas fluxes with open- and closed-path eddy covariance system: The WPL terms and spectral attenuation. In: X. Lee (Editor), Handbook of Micrometeorology: A guide for surface Flux Measurements. Kluwer, Dordrecht, pp. 133-160.
- *25 Fischer, G (Editor), 1988. Landolt-Börnstein, Numerical data and functional relationships in science and technology, Group V: Geophysics and space research, Volume 4: Meteorology Subvolume b: Physical and chemical properties of the air. Springer, Berlin, Heidelberg, 570pp.
- *26 Stull, R.B., 1988. An Introduction to Boundary Layer meteorology. Kluwer Acad. Publ., Dordrecht, Boston, London, 666pp.
- *27 Cohen, E. R. and Taylor, B. N., 1986. The 1986 adjustment of the fundamental physical constants. International Council of Scientific Unions (ICSU), Committee on Data for Science and Technology (CODATA). CODATA-Bulletin, No. 63: 36pp.

Quality control

- *28 Vickers, D. and Mahrt, L., 1997. Journal of Atmospheric and Oceanic Technology, 14: 512-526.
- *29 Foken, T. and Wichura, B., 1996. Agricultural and Forest Meteorology, 78: 83-105.
- *30 Hojstrup, J., 1993. Measuring Science Technology, 4: 153-157.
- *31 Schmid, H. P., 1994. Boundary-Layer Meteorology, 67: 293-318.
- *32 Korman, R. and Meixner, F.X., 1990. . Boundary-Layer Meteorology, 99: 207-224.