

1. About the data set

Site name (AsiaFlux three letter code)	Sapporo forest meteorology research site (SAP)	
Period of registered data	From January 1, 2001 to December 31, 2001	
This document file name	SAP_2001_001f.pdf	
Corresponding data file name	SAP_2001_001.csv SAP_2001P_001.csv (Hourly precipitation data)	
Revision information		
Date	Details of revision	Renewed file name
13 April 2010	First registration	SAP_2001_001.pdf SAP_2001_001.csv SAP_2001P_001.csv
8 June 2010	Document file is updated: p.9, line12: Publication list URL.	SAP_2001_002.pdf
21 July 2010	Document file is updated: p.9, line10-12: Publication information (title) is corrected.	SAP_2001_003.pdf
28 July 2010	Document file is renamed (no change in document contents): Version management is changed: 003 → 001c (corresponding data version + alphabetical sequence)	SAP_2001_001c.pdf
17 August 2011	Document file is updated: p.2, 13th row on right: Name of a dominant species is corrected.	SAP_2001_001d.pdf
15 May 2012	Document file is updated: p.9, line10: Publication information is updated.	SAP_2001_001e.pdf
31 July 2012	Document file is updated: New format is applied. p.1: Details in each revision are added. p.2: Dominant Species (Understory): Sasa Bamboo is corrected to Dwarf bamboo.	SAP_2001_001f.pdf
Contact person#1	[Flux & Meteorology] Katsumi YAMANOI (yamanoi@ffpri.affrc.go.jp)	
Contact person#2	[Flux & Meteorology] Kenzo KITAMURA (kitamu93@ffpri.affrc.go.jp)	
Contact person#3	[Flux & Meteorology] Yuichiro NAKAI (nakaiyui@ffpri.affrc.go.jp)	
Contact person#4	[Flux & Meteorology] Yasuko MIZOGUCHI (pop128@ffpri.affrc.go.jp)	

2. Site description

Hour line (Time difference from UTC)	Japan Standard Time (JST) (9 hours ahead of UTC)
Location (address)	Sapporo, Hokkaido, Japan
Position	42.9868N, 141.3853E (World Geodetic System 1984, GPS: Garmin eTrex Legend and map)
Elevation	182m above sea level (World Geodetic System 1984, GPS: Garmin eTrex Legend and map)
Terrain type	Gentle slope
Slope	0-7 degrees
Area	130 ha
Fetch	500-1500m
Climate	Cool temperate (Köppen climate classification: Dfb)
Mean annual air temperature	7.0 degree C (Tower site, 4years (2000-2003) average)
Mean annual precipitation	980 mm (Meteorological station(*1 *2), 4years (2000-2003) average)
Vegetation Type	Deciduous broadleaf forest
Dominant Species (Overstory)	Japanese white birch (<i>Betula platyphylla</i>), Mizunara oak (<i>Quercus crispula</i>), painted maple (<i>Acer mono</i>), aralia (<i>Kalopanax pictus</i>), Japanese linden (<i>Tilia japonica</i>), Japanese elm (<i>Ulmus davidiana</i> var. <i>japonica</i>)
Dominant Species (Understory)	Dwarf bamboo (<i>Sasa senanensis</i> , <i>Sasa kurilensis</i>)
Canopy height	Approx. 20m
Breast high diameter	approx. 25 cm (Japanese White Birch)
Age	90years (in 2002)
LAI	approx. 4 in max.
Soil Type	Black soil Bl _D (Andosol: WRB classification)
Other information	

Reference

*1 Hokkaido Research Center, Forestry and Forest Products Research Institute

FFPRI-Hokkaido (1998) Hitsujigaoka experimental forest operation plan. FFPRI-Hokkaido, Sapporo, Japan. pp1-44

*2 Sameshima R, Hirota T, Hamasaki T, Kato K, Iwata Y (2009) Meteorological Observation System at the National Agricultural Research Center for Hokkaido Region since 1966. Miscellaneous Publication of the National Agricultural Research Center for Hokkaido Region No.67. pp 1-8

3 Registered Data

Observation items	Symbol	Unit	Height(s) Depth(s)	Instruments	Note
Date	DATE	-	-	-	yyyymmdd
Time	TIME	-	-	-	hhmm
Precipitation	PPT	mm	1.8m		
Air temperature	Ta	degrees C	41.1m	HMP45D (VAISALA)	
Relative humidity	Rh	%	41.1m	HMP45D (VAISALA)	
Wind speed	U	$\text{m}\cdot\text{s}^{-1}$	41.4m	WM-30P (IKEDA)	
Wind direction	WD	degrees	28.5m	DA600-3T (KAIJO)	
Global solar radiation (incoming / downward)	Sd	$\text{W}\cdot\text{m}^{-2}$	41.3m	CM-6F (Kipp & Zonen)	
Reflected solar radiation (upward)	Su	$\text{W}\cdot\text{m}^{-2}$	39.0m	CM-6B (Kipp & Zonen)	
Photosynthetic active photon flux density (downward)	Pd	$\text{micromol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	41.2m	LI190 (LI-COR)	
Reflected PAR (upward)	Pu	$\text{micromol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	39.0m	LI190 (LI-COR)	
Net radiation	Rn	$\text{W}\cdot\text{m}^{-2}$	41.3m / 39.0m	CM-6F and CM-6B (Kipp & Zonen), PIR (Eppley)	4 elements (Sd, Su, Longwave radiation_down ward and Longwave radiation_upwa rd)
Soil heat flux	G	$\text{W}\cdot\text{m}^{-2}$	-0.02m	MF-81 (EKO)	
Sensible heat flux	H	$\text{W}\cdot\text{m}^{-2}$	28.5m	DA600-3T (KAIJO)	
Latent heat flux	IE	NA	NA	NA	
Friction velocity	Ust	$\text{m}\cdot\text{s}^{-1}$	28.5m	DA600-3T (KAIJO)	
CO ₂ flux	Fc	$\text{micromol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	28.5m	DA600-3T (KAIJO) LI-6262 (LI-COR)	Closed-path system QC
Storage change in canopy air layer	Sc	$\text{micromol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	2.7, 11.0, 16.7, 20.5, 30.0 m	LI-6262 (LI-COR)	
Net ecosystem exchange	NEE	$\text{micromol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	-	-	NEE=Fc+Sc Ust selection (Ust >= 0.38), another screenings *1 Gap filled *2

Ecosystem respiration	Re	$\text{micromol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	-	-	Missed Re at night and daytime Re were estimated using an empirical model ($\text{Re} = f(T_s)$) *2
Gross primary production	GPP	$\text{micromol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	-	-	GPP=-NEE+Re Gap filled *2

Reference

*1 $|\varphi| > 10^\circ$, $|\psi| > 10^\circ$ (in 3D rotation: first(θ), second(φ), third(ψ))→ eliminated

*2 -NEE = GPP + RE (daytime), -NEE = RE (nighttime)

GPP = $\text{Agmax} \cdot \alpha \cdot \text{PAR} / (\text{Agmax} + \alpha \cdot \text{PAR})$: Parameters were derived everyday. (from April to November)

GPP = 0 (from December to next March)

Re = -NEE(nighttime) = $a \cdot \exp(b \cdot T_s)$: Parameters (a & b) were derived yearly.

Data format

Data consists of fixed length (8 digits) comma separated format. Missing data is labeled as "-9999.00"

Line 1: Symbol (Date, Time, PPT, Ta,)

Line 2: Unit (yyyymmdd, hhmm, mm, degC,)

"hhmm" shows intermediate time of averaging period.

i.e. "1215" labels half-hourly average (or sum) of data from 12:00 to 12:30

Line 3: Comment

Line 4: Data

:

Data Example

Date, Time, yyyymmdd, hhmm,	PPT, mm,	Ta, degC,	Rh, %,	U, ms-1,	WD, deg,	Sd, Wm-2,	Su, Wm-2,	Pd, (#1),	Pu, (#1),	
File= KWG_2000_001.CSV; Created: 20100326; Gap= -9999.0; (*1): micro-mol m-2 s-1										
20000101, 0015,	0.0,	3.34,	87.19,	1.58,	-9999.0,	0.1,	-9999.0,	0.1,	0.0,	
20000101, 0045,	0.0,	3.12,	88.14,	1.44,	-9999.0,	0.0,	-9999.0,	0.1,	0.0,	
20000101, 0115,	0.0,	2.36,	90.51,	1.15,	-9999.0,	-0.3,	-9999.0,	0.1,	0.0,	
20000101, 0145,	0.0,	2.14,	91.32,	0.83,	-9999.0,	0.0,	-9999.0,	0.1,	0.0,	
20000101, 0215,	0.0,	2.28,	88.96,	0.49,	-9999.0,	-0.3,	-9999.0,	0.1,	0.0,	
20000101, 0245,	0.0,	2.24,	89.82,	0.35,	-9999.0,	-0.2,	-9999.0,	0.2,	0.0,	
20000101, 0315,	0.0,	2.05,	89.49,	1.50,	-9999.0,	0.1,	-9999.0,	0.2,	-0.1,	
20000101, 0345,	0.0,	2.41,	87.25,	1.27,	-9999.0,	0.0,	-9999.0,	0.2,	0.0,	
20000101, 0415,	0.0,	2.31,	86.83,	1.12,	-9999.0,	-0.2,	-9999.0,	0.1,	0.0,	
20000101, 0445,	0.0,	2.84,	83.36,	0.54,	-9999.0,	-0.6,	-9999.0,	0.0,	0.0,	
20000101, 0515,	0.0,	2.53,	83.32,	1.23,	-9999.0,	0.2,	-9999.0,	0.2,	0.0,	
20000101, 0545,	0.0,	1.59,	87.54,	1.29,	-9999.0,	-0.6,	-9999.0,	0.0,	0.0,	
20000101, 0615,	0.0,	1.89,	85.13,	0.84,	-9999.0,	0.4,	-9999.0,	0.3,	0.0,	
20000101, 0645,	0.0,	1.77,	82.40,	0.83,	-9999.0,	3.5,	-9999.0,	8.5,	0.4,	
20000101, 0715,	0.0,	2.67,	76.83,	1.38,	-9999.0,	45.8,	-9999.0,	71.9,	11.6,	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	
.	.	.	.	.	.	.	.	.	.	

4 Observation and calculation

4-1. Flux observation system and data acquisition

Type of sonic anemometer	DA600-3T (Probe TR-61C) (KAIJO)
Type of IRGA	LI-6262 (LI-COR)
Sampling rate	5Hz
Averaging time	30min
Flux measurement height #1	28.5m
Zero-plane displacement	
Roughness length	
Calibration information	
Other information	

4-2. Flux calculation

Calculation methods		Note
Flow attenuation ^{*4-6}	Not applied	
Coordinate rotation ^{*1-3}	3-D rotation	
Lag removal ^{*2, 7, 8}	Automatic	

4-3. Flux corrections

Correction methods		Target flux	Note
Cross wind correction ^{*9, 10}		For sensible heat flux (H)	
Vapor correction		For sensible heat flux (H)	
High frequency loss	Band-pass covariance method ^{*12}	For latent heat flux (IE)	
	Experimental approach ^{*2}	CO ₂ flux (Fc)	
Low frequency loss (Detrending)	Linear detrend ^{*16}	sensible heat flux (H), friction velocity, (Ust) CO ₂ flux (Fc)	
WPL Correction ^{*17-21}		For latent heat flux (IE) CO ₂ flux (Fc)	
Others ^{*22-24}	Temperature dependency for latent heat: L Humidity dependency for specific heat: Cp Temperature dependency for air density Pressure dependency for air density		

4-4. Quality control ^{*25-26}

QC methods			Note
Raw data test ^{*25,26}	Spike test ^{*27}	Applied	
	Absolute limits	Applied	
	Absolute variance		
	Higher-moment statistics	skewness	Applied
		kurtosis	Applied
	Discontinuities	Harr mean test	Applied
		Harr variance test	Applied
Visual inspection		Applied	
Non steady state test ^{*25}		Not applied	
Absolute thresholds			
Others			

4-5. Storage term

Target storage		Note
CO ₂	From CO ₂ profile data (2.7, 11.0, 16.7, 20.5, 30.0 m) Sampling interval: 120 seconds at each height	

References

--

5. Important events

Date	Events

6. Publications relating to this site

WATANABE Tsutomu, NAKAI Yuichiro, KITAMURA Kenzo, UTSUGI Hajime, TOBITA Hiroyuki, ISHIZUKA Shigehiro (2005) Scaling energy and CO₂ fluxes from leaf to canopy using a Multilayered Implementation for Natural Canopy-Environment Relations (MINCER). *Phyton*, 45:353-360

NAKAI Yuichiro, KITAMURA Kenzo, SUZUKI Satoru, ABE Shin (2003) Year-long carbon dioxide exchange above a broadleaf deciduous forest in Sapporo, Northern Japan. *Tellus B*, 55(3):305-312

SUZUKI Satoru, ISHIZUKA Shigehiro, KITAMURA Kenzo, YAMANOI Katsumi, NAKAI Yuichiro (2006) Continuous estimation of winter carbon dioxide efflux from the snow surface in a deciduous broadleaf forest. *Journal of Geophysical Research*, 111:D17101

KITAMURA Kenzo, NAKAI Yuichiro, SUZUKI Satoru, OHTANI Yoshikazu, YAMANOI Katsumi, SAKAMOTO Tomoki (2012) Interannual variability of net ecosystem production for a broadleaf deciduous forest in Sapporo, northern Japan, *Journal of Forest Research*, 17(3):323-332.

Publication list: http://www2.ffpri.affrc.go.jp/labs/flux/paper_e.html[SAP]

References

Flux calculation

- *1 McMillen, R.T., 1988. Boundary-Layer Meteorology, 43: 231-245.
- *2 Aubinet M. et al., 2000. Advances in Ecological Research, 30: 113-175.
- *3 Wilczak, J.M., Oncley, S.P. and Stage, S.A., 2001. Boundary-Layer Meteorology, 99: 127-150.
- *4 Wyngaard, J. C. and Zhang, S. F., 1985. J. Atmos. Oceanic Tech., 2: 548-558.
- *5 Kaimal, J.C. et al., 1990. Boundary-Layer Meteorol., 53: 103-115.
- *6 Shimizu, T. et al., 1999. Boundary-Layer Meteorol., 64: 227-236.
- *7 Leuning, R. and Judd M.J., 1996. Global Change Biology, 2: 241-254.
- *8 Information from Li-Cor

Flux correction

- *9 Schotanus, P. et al., 1983. Boundary-Layer Meteorology, 26: 81-93.
- *10 Liu, H., Peters, G. and Foken, T., 2001. Boundary-Layer Meteorology, 100: 459-468.
- *11 Kaimal J.C. and Gaynor, J.E., 1991. Boundary-Layer Meteorology, 56: 401-410.
- *12 Watanabe et al., 2000. Boundary-Layer Meteorol. 96, 743-491.
- *13 Massman, W. J., 2000. Agric. For. Meteorol. 104, 185-198
- *14 Massman, W. J., 2001. Agric. For. Meteorol. 107, 247-251
- *15 Moore, C.J., 1986. Boundary-Layer Meteorology, 37: 17-35.
- *16 Moncrieff, J. et al., 2004. Averaging, detrending and filtering of eddy covariance time series. In: X. Lee (Editor), Handbook of Micrometeorology: A guide for surface Flux Measurements. Kluwer, Dordrecht, pp. 7-31.
- *17 Webb, E. K., Pearman, G.I. and Leuning, R., 1980. Quarterly Journal of the Royal Meteorological Society, 106: 85-100.
- *18 Fuehrer, P.L. and Friehe, C.A., 2002. Boundary-Layer Meteorology, 102: 415-457.
- *19 Liebethal, C. and Foken, T., 2003. Boundary-Layer Meteorology, 109: 99-106.
- *20 Leuning, R. 2004. Measurements of trace gas fluxes in the atmosphere using eddy covariance: WPL corrections revisited. In: X. Lee (Editor), Handbook of Micrometeorology: A guide for surface Flux Measurements. Kluwer, Dordrecht, pp. 119-132.
- *21 Massman, W. 2004. Concerning the measurement of atmospheric trace gas fluxes with open- and closed-path eddy covariance system: The WPL terms and spectral attenuation. In: X. Lee (Editor), Handbook of Micrometeorology: A guide for surface Flux Measurements. Kluwer, Dordrecht, pp. 133-160.
- *22 Fischer, G (Editor), 1988. Landolt-Börnstein, Numerical data and functional relationships in science and technology, Group V: Geophysics and space research, Volume 4: Meteorology Subvolume b: Physical and chemical properties of the air. Springer, Berlin, Heidelberg, 570pp.
- *23 Stull, R.B., 1988. An Introduction to Boundary Layer meteorology. Kluwer Acad. Publ., Dordrecht, Boston, London, 666pp.
- *24 Cohen, E. R. and Taylor, B. N., 1986. The 1986 adjustment of the fundamental physical constants. International Council of Scientific Unions (ICSU), Committee on Data for Science and Technology (CODATA). CODATA-Bulletin, No. 63: 36pp.

Quality control

- *25 Vickers, D. and Mahrt, L., 1997. Journal of Atmospheric and Oceanic Technology, 14: 512-526.
- *26 Foken, T. and Wichura, B., 1996. Agricultural and Forest Meteorology, 78: 83-105.
- *27 Hojstrup, J., 1993. Measuring Science Technology, 4: 153-157.
- *28 Schmid, H. P., 1994. Boundary-Layer Meteorology, 67: 293-318.
- *29 Korman, R. and Meixner, F.X., 1990. . Boundary-Layer Meteorology, 99: 207-224.